

---

**Cytotoxic and regulatory T-cell molecule, 18-287aa, Human, His tag, Insect cell**

**Cat.NO.: TP01826**

3th Edition

**Synonyms:**CRTAM, CD355

**Description:**CRTAM, also known as cytotoxic and regulatory T-cell molecule, is a member of the immunoglobulin superfamily that complies with the structural characteristics of the JAM family of proteins and is phylogenetically more closely related to nectin-like proteins. It is a molecule involved in epithelial cell adhesion. This protein interacts with CADM1 promotes natural killer (NK) cell cytotoxicity and interferon-gamma (IFN-gamma) secretion by CD8+ cells in vitro as well as NK cell-mediated rejection of tumors expressing CADM3 in vivo. Recombinant human CRTAM, fused to His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.

**Form:**Liquid. In Phosphate Buffered Saline (pH 7.4) containing 10% glycerol.

**Molecular Weight:**31.0kDa (278aa) 40-57KDa (SDS-PAGE under reducing conditions.)

**Sequences:**

SLTNHTETITVEEGQTLTLKCVTSRLKNSSLQWLTPSGFTIFLNEYPALKNSKYQLLHHSANQLSITVPNVTLQDEGV  
YKCLHYSDSVSTKEVKVIVLATPFKPILEASVIRKQNGEEHVLMCSTMRSKPPPQITWLLGNSMEVSGGTLHEFET  
DGKKCNTTSTLIHTYGKNSTVDCIIRHRGLQGRKL VAPFRFEDLVTD EETASDALERNLSLSSQDPQQPTSTVSVTE  
DSSTSEIDKEEKEQTTQDPDLTTEANPQYLGLARKKSGLEHHHHHH

**Purity:**> 95% by HPLC

**Concentration:**1mg/ml (determined by Absorbance at 280nm)

**Endotoxin Level:**<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.