

**DUSP6, 1-381aa, Human, His tag, E.coli****Cat.NO.: TP01942**

3th Edition

**Synonyms:**Dual specificity phosphatase 6 isoform a, MKP3, PYST1

**Description:**DUSP6 is a member of the dual specificity protein phosphatase subfamily. These phosphatases inactivate their target kinases by dephosphorylating both the phosphoserine/threonine and phosphotyrosine residues. They negatively regulate members of the mitogen-activated protein (MAP) kinase superfamily (MAPK/ERK, SAPK/JNK, p38), which are associated with cellular proliferation and differentiation. Different members of the family of dual specificity phosphatases show distinct substrate specificities for various MAP kinases, different tissue distribution and subcellular localization, and different modes of inducibility of their expression by extracellular stimuli. Recombinant human DUSP6 protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.

**Form:**Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.15M NaCl, 10% glycerol, 1mM DTT

**Molecular Weight:**44.4 kDa (401aa) confirmed by MALDI-TOF

**Sequences:**

MGSSHHHHHHSSGLVPRGSHMIDTLRPVPPFASEMAISKTVAWLNEQLELGNERLLLMDCRPQELYESSHIESAINV  
AIPGIMLRRLQKGNLPVRALFTRGEDRDRFTRRCGTDTVVLYDESSSDWNENTGGESVLGLLLKKLKDEGCRAFY  
EGGFSKFQAEFSLHCETNLDGSCSSSSPPLPVLGLGGLRISSDSSSDIESDLDRDPNSATDSDGSPLSNSQPSFPV  
EILPFLYLGC AKDSTNLDVLEEFGIKYILNVTPNLPNLFENAGEFKYKQIPISDHWSQNLSQFFPEAISFIDEARGKNC  
GVLVHCLAGISRSVTVTVA YLMQKLNLSMNDAYDIVKMKKSNISPNFNFMGQLLDFERTLGLSSPCDNRVPAQQLY  
FTTPSNQNVYQVDSLQST

**Purity:**> 95% by HPLC

**Concentration:**0.25 mg/ml (determined by Bradford assay)

**Endotoxin Level:**<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.