

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

ECSIT, 19-217aa, Human, His tag, E.coli

Cat.NO.: TP01961

3th Edition

Synonyms: Evolutionarily conserved signaling intermediate in Toll pathway, mitochondrial, SITPEC.

Description:ECSIT is a ubiquitously expressed protein that plays an important role as an adaptor protein in the cytosolic signal transduction cascade events triggered by Toll receptor activation. It was initially identified as a cytoplasmic protein interacting specifically with TNF receptor associated factor (TRAF)-6 in the TLR pathway. Knockdown of ECSIT results in severely impaired complex I assembly and disturbed mitochondrial function. Recombinant human ECSIT protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.

Form:Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.1M NaCl, 10% glycerol,1mM DTT

Molecular Weight: 24.6 kDa (222aa) confirmed by MALDI-TOF

Sequences:

MGSSHHHHHHSSGLVPRGSHMGSGTCGAALTGTSISQVPLPKDSTGAADPPQPHIVGIQSPDQQAALARHNPARP VFVEGPFSLWLRNKCVYYHILRADLLPPEEREVEETPEEWNLYYPMQLDLEYVRSGWDNYEFDINEVEEGPVFAM CMAGAHDQATMAKWIQGLQETNPTLAQIPVVFRLAGSTRELQTSSAGLEEPPLPEDHQEEDDNLQRQQQGQS

Purity:> 95% by HPLC

Concentration: 1 mg/ml

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1