

**EPO1L, 24-468aa, Human, His tag, E.coli**

**Cat.NO.: TP02036**

3th Edition

**Synonyms:**ERO1-like protein alpha, ERO1-alpha, ERO1A

**Description:**ERO1-like protein alpha, also known as EPO1L, is an essential oxidoreductase that oxidizes proteins and is required for the folding of immunoglobulins. EPO1L covalently binds with PDI (protein disulfide-isomerase) and together they produce disulfide bonds between proteins in the endoplasmic reticulum. EPO1L is stimulated by hypoxia, suggesting that it is regulated through the HIF (hypoxia inducible transcription factor) pathway. EPO1L is ubiquitously expressed at low levels but expressed at high levels in upper digestive tract and esophagus. Recombinant human EPO1L protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.

**Form:**Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 10% glycerol

**Molecular Weight:**54.4 kDa (468aa)

**Sequences:**

MGSSHHHHHHSSGLVPRGSHMGSEEQPPETAARQCFCQVSGYLDDCTCDVETIDRFNNYRLFPRRLQKLLESDYF  
RYYKVNLRPCPFWNDISQCGRRDCAVKPCQSDEVPDGIKSASYKYSEEANNLIEECEQAERLGAVDESLSEETQ  
KAVLQWTKHDDSSDNFCEADDIQSPEAEYVDLLLNPERYTGYKGPDAWKIWNVIYEENCFKPQTIKRPLNPLASGQ  
GTSEENTFYSWLEGLCVEKRAFYRLISGLHASINVHLSARYLLQETWLEKKWGHNITEFQQRFDGILTEGEGPRRLK  
NLYFLYLIELRALSKVLPFFERPDFQLFTGNKIQDEENKMLLLEILHEIKSFPLHFDENSFFAGDKKEAHKLKEDFRLH  
FRNISRIMDCVGCFCRLWGKLQTQGLGTALKILFSEKLIANMPESGPSYEFHLTRQEIVSLFNAFGRISTSVKELEN  
FRNLLQNIH

**Purity:**> 95% by HPLC

**Concentration:**1 mg/ml (determined by Bradford assay)

**Endotoxin Level:**<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.