

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Erythropoietin, 28-193aa, Human, His tag, Insect cell

Cat.NO.: TP02049

3th Edition

Synonyms: EPO, EP, MVCD2

Description:EPO, also known as erythropoietin, is a glycoprotein hormone in the type I cytokine family and is related to thrombopoietin. Its three N-glycosylation sites, four alpha helices, and N- to C-terminal disulfide bond are conserved across species. It is primarily produced in the kidney by a population of fibroblast-like cortical interstitial cells adjacent to the proximal tubules. This protein can be found in the plasma and regulates red cell production by promoting erythroid differentiation and initiating hemoglobin synthesis. It also has neuroprotective activity against a variety of potential brain injuries and anti-apoptotic functions in several tissue types. Recombinant human EPO, fused to His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.

Form:Liquid. In Phosphate Buffered Saline (pH 7.4) containing 10% glycerol.

Molecular Weight:19.5kDa (174aa) 28-40KDa (SDS-PAGE under reducing conditions.)

Sequences:

APPRLICDSRVLERYLLEAKEAENITTGCAEHCSLNENITVPDTKVNFYAWKRMEVGQQAVEVWQGLALLSEAVLR GQALLVNSSQPWEPLQLHVDKAVSGLRSLTTLLRALRAQKEAISPPDAASAAPLRTITADTFRKLFRVYSNFLRGKL KLYTGEACRTGDRLEHHHHHH

Purity:> 95% by HPLC

Concentration:0.5mg/ml (determined by Absorbance at 280nm)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1