

F2, 328-622aa, Human, E.coli

Cat.NO.: TP02066

3th Edition

Synonyms:Coagulation factor II, Factor II (thrombin), PT

Description:Coagulation factor II, also known as F2, is proteolytically cleaved to form thrombin in the first step of the coagulation cascade which ultimately results in the stemming of blood loss. F2 also plays a role in maintaining vascular integrity during development and postnatal life. Mutations in F2 lead to various forms of thrombosis and dysprothrombinemia. Recombinant human F2 protein was expressed in E.coli.

Form:Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.4M Urea, 10% glycerol.

Molecular Weight:33.9kDa (296aa)

Sequences:

MTFGSGEADCGLRPLFEKKSLEDKTERELLESYIDGRIVEGSDAEIGMSPWQVMLFRKSPQELLCGASLISDRWWL
TAAHCLLYPPWDKNFTENDLLVRIGKHSRTRYERNIEKISMLEKIYIHPRYNWRENLRDIALMKLKKPVAFSDYIHPV
CLPDRETAASLLQAGYKGRVTGWGNLKETWTANVGKGQPSVLQVVNLPIVERPVCKDSTRITDNMFCAGYKPD
EGKRGDACEGDSGGPFVMKSPFNRRWYQMGIVSWGEGCDRDGKYGFYTHVFRLKKWIKVIDQFGE

Purity:> 95% by HPLC

Concentration:1.0 mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.