

## Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

FAAH2, 32-532aa, Human, His tag, E.coli

Cat.NO.: TP02069

3th Edition

Synonyms: Fatty-acid amide hydrolase 2, Fatty acid amide hydrolase 2, AMDD

**Description:**FAAH2 is a fatty acid amide hydrolase that shares a conserved protein motif with the amidase signature family of enzymes. The encoded enzyme is able to catalyze the hydrolysis of a broad range of bioactive lipids, including those from the three main classes of fatty acid amides; N-acylethanolamines, fatty acid primary amides and N-acyl amino acids. This enzyme has a preference for monounsaturated acyl chains as a substrate. Recombinant human FAAH2 protein, fused to His-tag at N-terminus, was expressed in E.coli .

Form:Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing, 10% glycerol, 0.4M Urea

Molecular Weight: 57.4kDa (524aa)

## Sequences:

MGSSHHHHHHSSGLVPRGSHMGSGGPKFASKTPRPVTEPLLLLSGMQLAKLIRQRKVKCIDVVQAYINRIKDVNPM INGIVKYRFEEAMKEAHAVDQKLAEKQEDEATLENKWPFLGVPLTVKEAFQLQGMPNSSGLMNRRDAIAKTDATVV ALLKGAGAIPLGITNCSELCMWYESSNKIYGRSNNPYDLQHIVGGSSGGEGCTLAAACSVIGVGSDIGGSIRMPAFF NGIFGHKPSPGVVPNKGQFPLAVGAQELFLCTGPMCRYAEDLAPMLKVMAGPGIKRLKLDTKVHLKDLKFYWMEH DGGSFLMSKVDQDLIMTQKKVVVHLETILGASVQHVKLKKMKYSFQLWIAMMSAKGHDGKEPVKFVDLLGDHGKH VSPLWELIKWCLGLSVYTIPSIGLALLEEKLRYSNEKYQKFKAVEESLRKELVDMLGDDGVFLYPSHPTVAPKHHVPL TRPFNFAYTGVFSALGLPVTQCPLGLNAKGLPLGIQVVAGPFNDHLTLAVAQYLEKTFGGWVCPGKF

Purity:> 95% by HPLC

Concentration: 1.0 mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1