

FEN1, 1-380aa, Human, E.coli

Cat.NO.: TP02110

3th Edition

Synonyms: Flap structure-specific endonuclease 1, FEN-1, MF1, RAD2.

Description: FEN-1, also known as Flap structure-specific endonuclease 1, removes 5' overhanging flaps in DNA repair and processes the 5' ends of Okazaki fragments in lagging strand DNA synthesis. The protein is a member of the XPG/RAD2 endonuclease family and is one of ten proteins essential for cell-free DNA replication. FEN-1 is highly homologous to yeast Rad2.

Form: Liquid. In 20 mM Tris-HCl buffer (pH8.0) containing 1mM DTT 10% glycerol

Molecular Weight: 42.5 kDa (380aa)

Sequences:

MGIQGLAKLIADVAPSAIRENDIKSYFGRKVAIDASMSIYQFLIAVRQGGDVLQNEEGETTSHLMGMFYRTIRMMEN
GIKPVYVFDGKPPQLKSGELAKRSERRAEAEKQLQQAQAAGAEQEVEKFTKRLVKVTKQHNDECKHLLSLMGIPYL
DAPSEAEASCAALVKAGKVYAAATEDMDCLTFGSPVLMRHLTASEAKKLPIQEFHLSRILQELGLNQEQFVDLCILL
GSDYCESIRGIGPKRAVDLIQKHKSIEEIVRRLDPNKYPVPENWLHKEAHLFLEPEVLDPEVELKWSEPNEEELIK
FMCGEKQFSEERIRSGVKRLSKSRQGSTQGRLLDDFFKVTGSLSSAKRKEPEPKGSTKKKAKTGAAGKFKRGK

Purity: > 95% by HPLC

Concentration: 1 mg/ml (determined by Bradford assay)

Endotoxin Level: <1.0 EU per 1 ug of protein (determined by LAL method)

Storage: Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.