

---

**Ferric uptake regulator(FUR), 1-148 aa E.coli, Recombinant, E.coli**

**Cat.NO.: TP02112**

3th Edition

**Synonyms:**DNA-binding transcriptional dual regulator of siderophore biosynthesis and transport.

**Description:**Fur (ferric uptake regulator) protein is a DNA-binding protein which regulates iron-responsive genes. Fur is a small, 17-kDa, global transcriptional repressor that in the presence of iron regulates functions as diverse as iron acquisition, oxidative stress, and virulence. In Escherichia coli, members of the Fur family regulate the expression of more than 100 genes that function in processes as varied as the biosynthesis and transport of siderophores, the expression of virulence factors, the alleviation of oxidative and NO-induced stress, and the inhibition of ferritin production through the expression of RyhB.

**Form:**Liquid. In 20 mM Tris-HCl buffer (pH 8.0) containing 2 mM CaCl<sub>2</sub>, 100 mM NaCl

**Molecular Weight:**16.7kDa (148aa), confirmed by MALDI-TOF

**Sequences:**

MTDNNTALKKAGLKVTLPRLKILEVLQEPDNHHVSAEDLYKRLIDMGEEIGLATVYRVLNQFDDAGIVTRHNFEGGK  
SVFELTQQHHHDHLICLDCGKVIEFSDDSIARQREIAAKHGIRLTNHSLYLYGHCAEGDCREDEHAHEGK

**Purity:**> 95% by HPLC

**Concentration:**1 mg/ml (determined by Bradford assay)

**Endotoxin Level:**<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.