

gldA, 1-367aa, E.coli, His tag, E.coli

Cat.NO.: TP02249

3th Edition

Synonyms:

Description:gldA catalyzes the NAD-dependent oxidation of glycerol to dihydroxyacetone (glycerone). This protein allows microorganisms to utilize glycerol as a source of carbon under anaerobic conditions. In E.coli, an important role of GldA is also likely to regulate the intracellular level of dihydroxyacetone by catalyzing the reverse reaction, i.e. the conversion of dihydroxyacetone into glycerol. gldA possesses a broad substrate specificity, since it is also able to oxidize 1,2-propanediol and to reduce glycolaldehyde, methylglyoxal and hydroxyacetone into ethylene glycol, lactaldehyde and 1,2-propanediol, respectively. Recombinant E. coli gldA protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.

Form:Liquid. In Phosphate buffered saline (pH7.4), 10% glycerol

Molecular Weight:41.1 kDa (390aa), confirmed by MALDI-TOF

Sequences:

MGSSHHHHHHSSGLVPRGSHMGSMRRIQSPGKYIQGADVINRLGEYLNKPLAERWLVGDKFVLGFAQSTVEKSF
KDAGLVVEIAPFGGECNQNEIDRLRGIAETAQCGAILGIGGGKTLDTAKALAHFMGVPVAIAPTIASTDAPCSALSVIY
TDEGEFDRYLLLPNNPNMVIVDTKIVAGAPARLLAAGIGDALATWFEARACSRSGATTMAGGKCTQAALALAEALCY
NTLLEEGEKAMLAEEQHVVTPALERVIEANTYLSGVGFESGGLAAAHAVHNGLTAIPTDAHYYHGEKVAFGTLTQLV
LENAPVEEIIETVAALSHAVGLPITLAQLDIKEDVPAKMRIVAEAAACAEGETIHNMPGGATPDQVYAALLVADQYGQRF
LQWE

Purity:> 95% by HPLC

Concentration:1 mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.