

GNG4, 1-72aa, Human, His tag, E.coli

## Cat.NO.: TP02295

**3th Edition** 

**Synonyms:** Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-4 precursor, Guanine nucleotide binding protein (G protein), gamma 4

**Description:**Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-4 precursor, also known as GNG4, is members of a multigene family and are implicated in determining the specificity of receptor-G protein interaction. In mammals, G protein alpha, beta and gamma polypeptides are encoded by at least 16, 4 and 7 genes, respectively. GNG4 is becoming increasingly clear that different G protein complexes expressed in different tissues carry structurally distinct members of the gamma as well as the alpha and beta subunits and that preferential association between members of subunit families increase G protein functional diversity. Recombinant human GNG4 protein, fused to His-tag at N-terminus, was expressed in E.coli.

Form:Liquid. In 20mM Tris-HCI buffer (pH 8.0) containing 0.4M Urea, 10% glycerol

Molecular Weight: 10.4 kDa (95aa)

## Sequences:

MGSSHHHHHHSSGLVPRGSHMGSMKEGMSNNSTTSISQARKAVEQLKMEACMDRVKVSQAAADLLAYCEAHVR EDPLIIPVPASENPFREKKFFC

Purity:> 95% by HPLC

Concentration: 0.5 mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.