

GREM2, 22-168aa, Human, His tag, E.coli

Cat.NO.: TP02333

3th Edition

Synonyms: Gremlin-2 precursor , CKTSF1B2, DAND3, PRDC

Description: GREM2 is a member of the BMP (bone morphogenic protein) antagonist family. Like BMPs, BMP antagonists contain cystine knots and typically form homo- and heterodimers. The CAN (cerberus and dan) subfamily of BMP antagonists, to which this protein belongs, is characterized by a C-terminal cystine knot with an eight-membered ring. The antagonistic effect of the secreted glycosylated protein is likely due to its direct binding to BMP proteins. As an antagonist of BMP, this protein may play a role in regulating organogenesis, body patterning, and tissue differentiation. Recombinant human GREM2 protein, fused to His-tag at N-terminus, was expressed in E.coli.

Form: Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.4M Urea, 10% glycerol

Molecular Weight: 19 kDa (170aa)

Sequences:

MGSSHHHHHSSGLVPRGSHMGSRKNRPAGAIPSPYKDGSSNNSERWQHQIKEVLASSQEALVVTERKYLKSDW
CKTQPLRQTVSEEGCRSRTILNRFYCYGQCNSFYIPRHVKKEEESFQSCAFCKPQRVTSVLVELECPGLDPPFRLKKI
QKVKQCRCMSVNLSDSDKQ

Purity: > 95% by HPLC

Concentration: 1 mg/ml (determined by Bradford assay)

Endotoxin Level: <1.0 EU per 1 ug of protein (determined by LAL method)

Storage: Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.