

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

GSTO1, 1-241aa, Human, E.coli

Cat.NO.: TP02360

3th Edition

Synonyms: Glutathione S-transferase omega-1, GSTTLp28, P28

Description:GSTO1, also known as p28 or GSTTLp28, is a protein that localizes to the cytoplasm and contains both an N-terminal and a C-terminal GST domain. In mouse, the encoded protein acts as a small stress response protein, likely involved in cellular redox homeostasis. This protein has dehydroascorbate reductase activity and may function in the glutathione-ascorbate cycle as part of antioxidant metabolism.

Form:Liquid. In 20 mM Tris-HCl buffer (pH8.0) containing 1mM DTT 10% glycerol

Molecular Weight: 27.5 kDa (241aa)

Sequences:

MSGESARSLGKGSAPPGPVPEGSIRIYSMRFCPFAERTRLVLKAKGIRHEVININLKNKPEWFFKKNPFGLVPVLEN SQGQLIYESAITCEYLDEAYPGKKLLPDDPYEKACQKMILELFSKVPSLVGSFIRSQNKEDYAGLKEEFRKEFTKLEE VLTNKKTTFFGGNSISMIDYLIWPWFERLEAMKLNECVDHTPKLKLWMAAMKEDPTVSALLTSEKDWQGFLELYLQ NSPEACDYGL

Purity:> 95% by HPLC

Concentration: 1 mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1