

## Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

HAO1, 1-370aa, Human, His tag, E.coli

Cat.NO.: TP02398

3th Edition

Synonyms: Hydroxyacid oxidase 1, GOX (glycolate oxidase), GOX1, HAOX1.

**Description:**HAO1, also known as glycolate oxidase, is a member of the superfamily of the alpha hydroxy acid oxidases(HAO) enzymes. It catalyzes the FMN mediated oxidation of glycolate to glyoxylate and glyoxylate to oxalate with reduction of oxygen to hydrogen peroxide. It is most highly expressed in liver and pancreas and is most active on twocarbon substrates such as glycolate.

Form:Liquid. 20mM Tris-HCl buffer (pH8.0) containing 20% glycerol 0.5M NaCl

Molecular Weight: 45.0 kDa(406aa), confirmed by MALDI-TOF

## Sequences:

MRGSHHHHHHGMASMTGGQQMGRDLYDDDDKDRWGSMLPRLICINDYEQHAKSVLPKSIYDYYRSGANDEETL ADNIAAFSRWKLYPRMLRNVAETDLSTSVLGQRVSMPICVGATAMQRMAHVDGELATVRACQSLGTGMMLSSWA TSSIEEVAEAGPEALRWLQLYIYKDREVTKKLVRQAEKMGYKAIFVTVDTPYLGNRLDDVRNRFKLPPQLRMKNFET STLSFSPEENFGDDSGLAAYVAKAIDPSISWEDIKWLRRLTSLPIVAKGILRGDDAREAVKHGLNGILVSNHGARQLD GVPATIDVLPEIVEAVEGKVEVFLDGGVRKGTDVLKALALGAKAVFVGRPIVWGLAFQGEKGVQDVLEILKEEFRLA MALSGCQNVKVIDKTLVRKNPLAVSKI

Purity:> 95% by HPLC

Concentration: 1.0 mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1