
HCV-E2, 482-671aa, HCV, His tag, E.coli**Cat.NO.: TP02417**

3th Edition

Synonyms: Polyprotein,**Description:** E1 and E2 glycoproteins form a heterodimer that is involved in virus attachment to the host cell, virion internalization through clathrin-dependent endocytosis and fusion with host membrane. E1/E2 heterodimer binds to human LDLR, CD81 and SCARB1/SR-BI receptors, but this binding is not sufficient for infection, some additional liver specific cofactors may be needed. The fusion function may possibly be carried by E1. E2 inhibits human EIF2AK2/PKR activation, preventing the establishment of an antiviral state. E2 is a viral ligand for CD209/DC-SIGN and CLEC4M/DC-SIGNR, which are respectively found on dendritic cells (DCs), and on liver sinusoidal endothelial cells and macrophage-like cells of lymph node sinuses. These interactions allow capture of circulating HCV particles by these cells and subsequent transmission to permissive cells. Recombinant HCV (Hepatitis C Virus) E2 protein, fused to His-tag at N-terminus, was expressed in E.coli.**Form:** Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.4M Urea, 10% glycerol**Molecular Weight:** 25.4 kDa (226aa)**Sequences:**

```
MRGSHHHHHHGMASMTGGQMQGRDLYDDDDKDRWGSERPWCWHYPPRPGIVPAKSVCGPVYCFTPSPVVVG  
TTDRSGAPTYSWGANDTDVFLNTRPPLGNWFGCTWMNSTGFTKVCGAPPCVIGGVGNNTLLCPTDCFRKHPE  
ATYSRCGSGPWITPRCMVDYPYRLWHYPCTINYTIFKVRMYVGGVEHRLEAACNWTRGERCDLEDRDRSELSPLL  
LSTTQ
```

Purity: > 95% by HPLC**Concentration:** 0.25 mg/ml (determined by Bradford assay)**Endotoxin Level:** <1.0 EU per 1 ug of protein (determined by LAL method)**Storage:** Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.