

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

HEXA, 23-529aa, Human, His tag, Insect cell

Cat.NO.: TP02438

3th Edition

Synonyms: Beta-hexosaminidase subunit alpha preproprotein, TSD

Description:HEXA, also known as beta-hexosaminidase subunit alpha preproprotein, is an enzyme, located on the 15th chromosome. Hexosaminidase A and the cofactor GM2 activator protein catalyze the degradation of the GM2 gangliosides and other molecules containing terminal N-acetyl hexosamines. Hexosaminidase A is a heterodimer composed of an alpha subunit and a beta subunit. Recombinant human HEXA, fused to His-tag at C-terminus, was expressed in insect cell and purified by using conventional chromatography techniques.

Form:Liquid. In phosphate buffered saline (pH7.4).

Molecular Weight: 59.2kDa (513aa) 50-70kDa (SDS-PAGE under reducing conditions.)

Sequences:

LWPWPQNFQTSDQRYVLYPNNFQFQYDVSSAAQPGCSVLDEAFQRYRDLLFGSGSWPRPYLTGKRHTLEKNVLV VSVVTPGCNQLPTLESVENYTLTINDDQCLLLSETVWGALRGLETFSQLVWKSAEGTFFINKTEIEDFPRFPHRGLLL DTSRHYLPLSSILDTLDVMAYNKLNVFHWHLVDDPSFPYESFTFPELMRKGSYNPVTHIYTAQDVKEVIEYARLRGIR VLAEFDTPGHTLSWGPGIPGLLTPCYSGSEPSGTFGPVNPSLNNTYEFMSTFFLEVSSVFPDFYLHLGGDEVDFTC WKSNPEIQDFMRKKGFGEDFKQLESFYIQTLLDIVSSYGKGYVVWQEVFDNKVKIQPDTIIQVWREDIPVNYMKELE LVTKAGFRALLSAPWYLNRISYGPDWKDFYIVEPLAFEGTPEQKALVIGGEACMWGEYVDNTNLVPRLWPRAGAV AERLWSNKLTSDLTFAYERLSHFRCELLRRGVQAQPLNVGFCEQEFEQTHHHHHHH

Purity:> 95% by HPLC

Concentration: 0.5 mg/ml (determined by Absorbance at 280 nm)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1