

**HLA-DOB, 27-224aa, Human, His tag, E.coli**

**Cat.NO.: TP02460**

3th Edition

**Synonyms:**Major histocompatibility complex, class II, DO beta, DOB

**Description:**HLA-DOB belongs to the HLA class II beta chain paralogues. This class II molecule is a heterodimer consisting of an alpha (DOA) and a beta chain (DOB), both anchored in the membrane. It is located in intracellular vesicles. DO suppresses peptide loading of MHC class II molecules by inhibiting HLA-DM. Class II molecules are expressed in antigen presenting cells (APC: B lymphocytes, dendritic cells, macrophages). The beta chain is approximately 26-28 kDa and its gene contains 6 exons. Exon one encodes the leader peptide, exons 2 and 3 encode the two extracellular domains, exon 4 encodes the transmembrane domain and exon 5 encodes the cytoplasmic tail. Recombinant human HLA-DOB protein, fused to His-tag at N-terminus, was expressed in E.coli.

**Form:**Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.4M Urea, 10% glycerol

**Molecular Weight:**25.2 kDa (222aa)

**Sequences:**

MGSSHHHHHHSSGLVPRGSHMGSGTDSPEDFVIQAKADCYFTNGTEKVQFVVRFIFNLEEYVRFDSDDVGMFVALT  
KLGQPDAEQWNSRLDLLERSRQAVDGVCRHNYRLGAPFTVGRKVQPEVTVPERTPLLHQHNLLHCSVTGFYPG  
DIKIKWFLNGQEERAGVMSTGPIRNGDWTFQTVVMLEMTPELGHVYTCLVDHSSLLSPVSVEWRAQSEYSWRK

**Purity:**> 95% by HPLC

**Concentration:**1 mg/ml (determined by Bradford assay)

**Endotoxin Level:**<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.