

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

HP, 19-347aa Human, His tag, E.coli

Cat.NO.: TP02489

3th Edition

Synonyms: Haptoglobin isoform 2 preproprotein, BP, HP2ALPHA2, HPA1S

Description:HP, as known as haptoglobin, is processed to yield both alpha and beta chains, which subsequently combine as a tetramer to produce haptoglobin. In blood plasma, haptoglobin binds free hemoglobin (Hb) released from erythrocytes with high affinity and thereby inhibits its oxidative activity. The haptoglobin-hemoglobin complex will then be removed by the reticuloendothelial system (mostly the spleen). Recombinant human HP protein, fused to His-tag at N-terminus, was expressed in E.coli

Form:Liquid. 20mM Tris-HCl buffer (pH8.0) containing 10% glycerol 0.4M Urea

Molecular Weight: 39.0 kDa(352aa)

Sequences:

MGSSHHHHHHSSGLVPRGSHMGSVDSGNDVTDIADDGCPKPPEIAHGYVEHSVRYQCKNYYKLRTEGDGVYTLN NEKQWINKAVGDKLPECEAVCGKPKNPANPVQRILGGHLDAKGSFPWQAKMVSHHNLTTGATLINEQWLLTTAKN LFLNHSENATAKDIAPTLTLYVGKKQLVEIEKVVLHPNYSQVDIGLIKLKQKVSVNERVMPICLPSKDYAEVGRVGYV SGWGRNANFKFTDHLKYVMLPVADQDQCIRHYEGSTVPEKKTPKSPVGVQPILNEHTFCAGMSKYQEDTCYGDA GSAFAVHDLEEDTWYATGILSFDKSCAVAEYGVYVKVTSIQDWVQKTIAEN

Purity:> 95% by HPLC

Concentration: 0.5mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1