

HPRT, 1-218aa, Human, His-tag, E.coli (Bioactivity Validated)

Cat.NO.: TP02496

3th Edition

Synonyms:Hypoxanthine-guanine phosphoribosyltransferase, HGPRT, HGPRTase, HPRT

Description:Hypoxanthine-guanine phosphoribosyltransferase, also known as HPRT1 has a central role in the generation of purine nucleotides through the purine salvage pathway. The enzyme primarily functions to salvage purines from degraded DNA to renewed purine synthesis. In this role, it acts as a catalyst in the reaction between guanine and phosphoribosyl pyrophosphate to form GMP. Recombinant human HPRT1, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.

Form:Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing, 20% glycerol

Molecular Weight:26.7 kDa (238aa) confirmed by MALDI-TOF

Sequences:

MGSSHHHHHHSSGLVPRGSHMATRSPGVVISDDEPGYDLDFCIPNHYAEDLERVFIPHGLIMDRTERLARDVMKE
MGGHHIVALCVLKGGYKFFADLLDYIKALNRNSDRSIPMTVDFIRLKSVCNDQSTGDIKVI GGDDLSTLTGKNVLIVED
IIDTGKTMQTL LSLVRQYNPKMVKVASLLVKRTPRSVGYKPDFVGF EIPDKFVVG YALDYNEYFRDLNHVCV ISETG
KAKYKA

Purity:> 95% by HPLC

Concentration:0.5mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.