

IDH3G, 40-393aa, Human, His tag, E.coli

Cat.NO.: TP02554

3th Edition

Synonyms: Isocitrate dehydrogenase [NAD] subunit gamma, mitochondrial, H-IDHG.

Description: IDH3G, also known as isocitrate dehydrogenase [NAD] subunit gamma, mitochondrial, belongs to the isocitrate and isopropylmalate dehydrogenases family. Isocitrate dehydrogenases catalyze the oxidative decarboxylation of isocitrate to 2-oxoglutarate. IDH3G is a gamma subunit of one isozyme of isocitrate dehydrogenase that belongs to a distinct subclass, which utilizes NAD(+) as the electron acceptor, and is localised to the mitochondrial matrix.

Form: Liquid. 20mM Tris-HCl buffer (pH8.0) containing 50% glycerol, 0.2M NaCl, 5mM DTT, 2mM EDTA

Molecular Weight: 41.1 kDa (375aa), confirmed by MALDI-TOF

Sequences:

MGSSHHHHHHSSGLVPRGSHMFSEQTIPPSAKYGGRHTVTMIPGDGIGPELMLHVKS VFRHACVPVDFEEVHVSS
NADEEDIRNAIMAIRNRNRVALKGNIETNHNLP SHKSRNNILRTSLDLYANVIHCKSLPGVVTRHKDIDILIVRENTGE
YSSLEHESVAGVVESLK IITKAKSLRIA EYAFKLAQESGRKKVTAVHKANIMKLGDGLFLQCCREVAARYPQITFENMI
VDNTTMQLVSRPQQFDVMVMPNLYGNIVNNVCAGLVGGPGLVAGANYGHVYAVFETATRNTGKSIANKNIANPTA
TLLASCMMLDHLKLHSYATSIRKAVLASMDNENMHTPDIGGQGT TSEAIQDVIRHIRVINGRAVEA

Purity: > 95% by HPLC

Concentration: 0.25 mg/ml (determined by Bradford assay)

Endotoxin Level: <1.0 EU per 1 ug of protein (determined by LAL method)

Storage: Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.