

IGJ, 23-159aa, Human, His tag, E.coli

Cat.NO.: TP02587

3th Edition

Synonyms:Immunoglobulin J chain, IGCJ, JCH

Description:Immunoglobulin J chain, also known as IGJ, serves to link two monomer units of either IgM or IgA. In the case of IgM, the J chain-joined dimer is a nucleating unit for the IgM pentamer, and in the case of IgA it induces larger polymers. It also helps to bind these immunoglobulins to secretory component. Recombinant human IGJ protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.

Form:Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.15M NaCl, 10% glycerol, 1mM DTT

Molecular Weight:18.0 kDa (160aa) confirmed by MALDI-TOF

Sequences:

MGSSHHHHHSSGLVPRGSHMGSQEDERIVLVDNKCKCARITSRIIRSSDPNEDIVERNIRIIVPLNNRENISDPTS
PLRTRFVYHLSDLCKKCDPTEVELDNQIVTATQSNICDEDSATETCYTYDRNKCYTAVVPLVYGGETKMOVETALTPD
ACYPD

Purity:> 95% by HPLC

Concentration:0.5 mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.