

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

IL-33, 112-270aa, Human, Recombinant, E.coli

Cat.NO.: TP02599

3th Edition

Synonyms:Interleukin-33, DVS27, NF-HEV, NKHEV, C9orf26

Description:Interleukin-33 (IL-33) is a 32kDa proinflammatory cytokine that may also regulate gene transcription in producer cells. IL-33 is structurally related to IL-1, which induces helper T cells to produce type 2 cytokines and acts through the receptor IL1RL-1 (IL1 receptor-like-1), which is known also as ST2. Binding of IL-33 to this receptor activates NF-kappa-B and MAP kinases and induces in vitro Th2 cells to produce cytokines.

Form:Liquid. In 20 mM Tris-HCl buffer (pH 8.0)

Molecular Weight: 18.1Da (160aa), confirmed by MALDI-TOF

Sequences:

MSITGISPITEYLASLSTYNDQSITFALEDESYEIYVEDLKKDEKKDKVLLSYYESQHPSNESGDGVDGKMLMVTLSP TKDFWLHANNKEHSVELHKCEKPLPDQAFFVLHNMHSNCVSFECKTDPGVFIGVKDNHLALIKVDSSENLCTENILF KLSET

Purity:> 95% by HPLC

Concentration:1 mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1