

## Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

IL18BP, 31-194aa Human, His tag, E.coli

Cat.NO.: TP02620

3th Edition

Synonyms: Interleukin-18-binding protein, IL18BPa

**Description:**IL18BP functions as an inhibitor of the proinflammatory cytokine, IL18. It binds IL18, prevents the binding of IL18 to its receptor, and thus inhibits IL18-induced IFN-gamma production, resulting in reduced T-helper type 1 immune responses. This protein is constitutively expressed and secreted in mononuclear cells. Elevated level of this protein is detected in the intestinal tissues of patients with Crohn's disease. Recombinant human IL18BP protein, fused to His-tag at N-terminus, was expressed in E.coli.

Form:Liquid. In 20mM Tris-HCl buffer (pH 8.0) containing 0.4M UREA, 10% glycerol

Molecular Weight: 20.0 kDa (187aa)

## Sequences:

MGSSHHHHHHSSGLVPRGSHMGSTPVSQTTTAATASVRSTKDPCPSQPPVFPAAKQCPALEVTWPEVEVPLNGT LSLSCVACSRFPNFSILYWLGNGSFIEHLPGRLWEGSTSRERGSTGTQLCKALVLEQLTPALHSTNFSCVLVDPEQ VVQRHVVLAQLWAGLRATLPPTQEALPSSHSSPQQG

Purity:> 95% by HPLC

Concentration: 1 mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1