

IMPDH2, 1-514aa, Human, His tag, E.coli

Cat.NO.: TP02661

3th Edition

Synonyms: Inosine-5'-monophosphate dehydrogenase 2, IMPD2, IMPDH-II

Description:IMPDH2 belongs to the IMPDH/GMPR family. It catalyzes the NAD-dependent oxidation of inosine-5'-monophosphate into xanthine-5'-monophosphate, which is then converted into guanosine-5'-monophosphate. IMPDH2 is the rate-limiting enzyme in the de novo guanine nucleotide biosynthesis. It is thus involved in maintaining cellular guanine deoxy- and ribonucleotide pools needed for DNA and RNA synthesis. Also IMPDH1 and IMPDH2 are targets for the important immunosuppressive drug, mycophenolic acid (MPA). Recombinant human IMPDH2 protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography techniques.

Form:Liquid. In 20mM Tris-HCI buffer (pH 8.0) containing 2mM DTT, 20% glycerol, 150mM NaCI

Molecular Weight: 58.0kDa (534aa)

Sequences:

Purity:> 95% by HPLC

Concentration: 0.5mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.