

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

NME4, 33-187aa, Human, His tag, E.coli

Cat.NO.: TP03172

3th Edition

Synonyms: Nucleoside diphosphate kinase, mitochondrial, NDK, NDPKD, nm23-H4, NM23D

Description:NME4, also known as nucleoside diphosphate kinase, mitochondrial, belongs to the NDK family. NME4 are ubiquitous enzymes that catalyze transfer of gamma-phosphates, via a phosphohistidine intermediate, between nucleoside and dioxynucleoside tri- and diphosphates. The enzymes are products of the nm23 gene family, which includes NME4. NME4 plays a major role in the synthesis of nucleoside triphosphates other than ATP.

Form:Liquid. 20mM Tris-HCl buffer (pH8.0) containing 40% glycerol,0.2M NaCl

Molecular Weight: 19.6 kDa (176aa) confirmed by MALDI-TOF

Sequences:

MGSSHHHHHHSSGLVPRGSHMPSWTRERTLVAVKPDGVQRRLVGDVIQRFERRGFTLVGMKMLQAPESVLAEH YQDLRRKPFYPALIRYMSSGPVVAMVWEGYNVVRASRAMIGHTDSAEAAPGTIRGDFSVHISRNVIHASDSVEGAQ REIQLWFQSSELVSWADGGQHSSIHPA

Purity:> 95% by HPLC

Concentration: 0.5 mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1