

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

SNTN, 1-147aa, Human, His tag, E.coli

Cat.NO.: TP03994

3th Edition

Synonyms: Sentan, S100A1L, S100AL

Description:SNTN, also known as sentan, belongs to the S-100 family. SNTN is shown to localize exclusively to the bridging structure between the cell membrane and peripheral singlet microtubules, which specifically exists in the narrowed distal portion of cilia. Exogenously expressed sentan showed affinity for the membrane protrusions, and a protein-lipid binding assay revealed that sentan bound to phosphatidylserine. These findings suggest that sentan is the first molecular component of the ciliary tip to bridge the cell membrane and peripheral singlet microtubules, making the distal portion of the cilia narrow and stiff to allow for better airway clearance or ovum transport. Recombinant human SNTN protein, fused to His-tag at N-terminus, was expressed in E.coli and purified by using conventional chromatography.

Form:Liquid. 20mM Tris-HCl buffer (pH8.0) containing 50% glycerol, 0.15M NaCl, 1mM DTT

Molecular Weight: 18.6 kDa (167aa), confirmed by MALDI-TOF

Sequences:

MGSSHHHHHHSSGLVPRGSHMGGCMHSTQDKSLHLEGDPNPSAAPTSTCAPRKMPKRISISKQLASVKALRKCS DLEKAIATTALIFRNSSDSDGKLEKAIAKDLLQTQFRNFAEGQETKPKYREILSELDEHTENKLDFEDFMILLLSITVMS DLLQNIRNVKIMK

Purity:> 95% by HPLC

Concentration:0.5 mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1