

TANK, 1-119 aa, Human, Recombinant, E.coli

Cat.NO.: TP04142

3th Edition

Synonyms:TRAF interacting protein TANK isoform b, I-TRAF, TRAF2, TRAF interacting protein TANK isoform b I TRAF, ITRAF, TRAF interacting protein, TRAF family member associated NF KAPPA B activator, TRAF family member associated NFKB activator

Description:TANK, also known as TRAF(tumor necrosis factor receptor-associated Factor)-interacting protein, is found in the cytoplasm and can bind to TRAF1, TRAF2, or TRAF3, thereby proposing that this protein is an inhibitor of TRAF function that regulates TRAF protein activity by sequestering TRAFs in a latent state in the cytoplasm. Overexpression of TANK inhibits TRAF2-mediated NF-Kappa-B activation signaled by CD40 and both TNF receptors and inhibits LMP1-mediated NF-kappa-B activation by blocking the association of TRAF2 with LMP1. Recombinant human TANK protein was expressed in E.coli and purified by using conventional chromatography techniques.

Form:Liquid. In 20mM Tris-HCl buffer (pH8.0)

Molecular Weight:13.6 kDa (119 aa), confirmed by MALDI-TOF.

Sequences:

MDKNIGEQLNKAYEAFRQACMDRDSAVKELQQKTENYEQRIREQQEQLSLQQTIIDKLKSQLLLLVNSTQDNNYGCV
PLLEDSETRKNNLTDQPQDKVISGIAREKLPKVDIASAESSI

Purity:> 95% by HPLC

Concentration:1 mg/ml (determined by Bradford assay)

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.