

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Recombinant Human Tumor Necrosis Factor Receptor II/TNFRSF1B/CD120b Protein(C-Fc)

Cat.NO.: TP05075

3th Edition

Synonyms:Tumor necrosis factor receptor superfamily member 1B; TNFRSF1B; Tumor necrosis factor receptor 2; TNF-R2; TNF-RII; Tumor necrosis factor receptor type II; p75; p80 TNF-alpha receptor; CD120b

Description:Tumor necrosis factor receptor superfamily member 1B (TNFRSF1B) is a member of the tumor necrosis factor receptor superfamily. Human TNF RII contains four cysteinerich repeats in its ECD, which shares 58% and 56% amino acid sequence identity with the mouse and rat orthologs, respectively.TNF RII is expressed predominantly on cells of the hematopoietic lineage, such as T and natural killer cells, as well as on endothelial cells, microglia, astrocytes, neurons, oligodendrocytes, cardiac myocytes, thymocytes, and mesenchymal stem cells.TNF RII binds to the membranebound forms of TNF? and Lymphotoxin?/TNF??soluble TNF is thought to signal predominately through TNF RI.Soluble TNF RII is believed to inhibit TNF biological activity by binding TNF thereby preventing it from activating membrane TNF receptors.

Form:PBS

Molecular Weight:51.9 kDa

Sequences:Leu23-Asp257

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1