

Recombinant Mouse Cell adhesion molecule 1 Protein

Cat.NO.: TP05223

3th Edition

Synonyms:Cell adhesion molecule 1;Immunoglobulin superfamily member 4;IgSF4;Nectin-like protein 2;NECL-2;Synaptic cell adhesion molecule;SynCAM;Tumor suppressor in lung cancer 1;TSLC-1

Description:Cell adhesion molecule 1(CADM1) is a single-pass type I membrane protein and belongs to the nectin family. It contains 2 Ig-like C2-type (immunoglobulin-like) domains and 1 Ig-like V-type (immunoglobulin-like) domain. CADM1 acts as a tumor suppressor in non-small-cell lung cancer (NSCLC) cells. Interaction with CRTAM promotes natural killer (NK) cell cytotoxicity and interferon-gamma (IFN-gamma) secretion by CD8+ cells in vitro as well as NK cell-mediated rejection of tumors expressing CADM3 in vivo. CADM1 may contribute to the less invasive phenotypes of lepidic growth tumor cells. In mast cells, it may mediate attachment to and promote communication with nerves. CADM1, together with MITF, is essential for development and survival of mast cells in vivo. The protein acts as a synaptic cell adhesion molecule and plays a role in the formation of dendritic spines and in synapse assembly. It may be involved in neuronal migration, axon growth, pathfinding, and fasciculation on the axons of differentiating neurons. CADM1 may play diverse roles in the spermatogenesis including in the adhesion of spermatocytes and spermatids to Sertoli cells and for their normal differentiation into mature spermatozoa.

Form:PBS

Molecular Weight:65.3 kDa

Sequences:Met1-His377

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.