

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Recombinant Human Hepatocyte Growth Factor/HGF/Hepatopoietin-A Protein(C-6His)

Cat.NO.: TP05265

3th Edition

Synonyms: Hepatocyte growth factor; HPTA; HGF; SF; Scatter factor; Hepatopoietin-A

Description:Hepatocyte growth factor/scatter factor (HGF/SF) is a paracrine cellular growth, motility and morphogenic factor. It belongs to the peptidase S1 family and Plasminogen subfamily, contains 4 kringle domains, 1 PAN domain and 1 peptidase S1 domain. HGF regulates cell growth, cell motility, and morphogenesis by activating a tyrosine kinase signaling cascade after binding to the proto-oncogenic c-Met receptor. HGF is secreted by mesenchymal cells and acts as a multi-functional cytokine on cells of mainly epithelial origin. Its ability to stimulate mitogenesis, cell motility, and matrix invasion gives it a central role in angiogenesis, tumorogenesis, and tissue regeneration.

Form:PBS

Molecular Weight: 26&53.7&80.7 kDa

Sequences: Gln32-Ser728

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1