
Recombinant Mouse Interleukin-17F/IL-17F Protein(C-6His)**Cat.NO.: TP05335**

3th Edition

Synonyms:Interleukin-17F; IL-17F; Cytokine ML-1; Interleukin-24; IL-24; IL17F; IL24**Description:**Interleukin-17F (IL-17F) exists in a disulfide-linked heterodimer that belongs to the IL-17 family. IL-17F is expressed in activated, but not resting, CD4+ T-cells and activated monocytes. Mouse and human IL-17F share 55% sequence identity.IL-17F has been shown to stimulate the production of several other cytokines, including IL-6, IL-8, and granulocyte colony-stimulating factor. IL-17F can regulate cartilage matrix turnover and stimulates PBMC and T-cell proliferation. IL-17F is also found to inhibit the angiogenesis of endothelial cells and induce endothelial cells to produce IL2, TGFB1/TGFB, and monocyte chemoattractant protein-1. Defects in IL-17F are the cause of familial candidiasis type 6 (CANDF6).**Form:**PBS**Molecular Weight:**15.9 kDa**Sequences:**Arg29-Ala161**Purity:**> 95% by HPLC**Concentration:****Endotoxin Level:**<1.0 EU per 1 ug of protein (determined by LAL method)**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.