

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Recombinant Human Signal Transducer and Activator of Transcription 1 Protein

Cat.NO.: TP05351

3th Edition

Synonyms:Signal Transducer and Activator of Transcription 1-Alpha/Beta; Transcription Factor ISGF-3 Components p91/p84; STAT1

Description:Signal Transducer and Activator of Transcription 1-Alpha/Beta (STAT1) contains one SH2 domain and belongs to the transcription factor STAT family. When tyrosine- and serine-phosphorylated, STAT1 can form a homodimer termed IFN-gamma-activated factor (GAF), migrate into the nucleus and bind to the IFN gamma activated sequence (GAS) to drive the expression of the target genes, inducing a cellular antiviral state. STAT1 functions as signal transducer and transcription activator that mediates cellular responses to interferons. Defects in STAT1 are the cause of STAT1 deficiency complete and familial candidiasis type 7.

Form:PBS

Molecular Weight:83.3 kDa

Sequences:Met 1-Val712

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1