

**Recombinant Human Aldo-Keto Reductase 1C3/AKR1C3 Protein(C-6His)**

**Cat.NO.: TP05379**

3th Edition

**Synonyms:**Aldo-Keto Reductase Family 1 Member C3; 17-Beta-Hydroxysteroid Dehydrogenase Type 5; 17-Beta-HSD 5; 3-Alpha-HSD Type II Brain; 3-Alpha-Hydroxysteroid Dehydrogenase Type 2; 3-Alpha-HSD Type 2; Chlordecone Reductase Homolog HAKRb; Dihydrodiol Dehydrogenase 3; DD-3; DD3; Dihydrodiol Dehydrogenase Type I; HA1753; Indanol Dehydrogenase; Prostaglandin F Synthase; Testosterone 17-Beta-Dehydrogenase 5; Trans-1;2-Dihydrobenzene-1;2-Diol Dehydrogenase; AKR1C3; DDH1; HSD17B5; KIAA0119; PGFS

**Description:**AKR1C3, is an enzyme which belongs to the aldo/keto reductase family. It is expressed in many tissues including adrenal gland, brain, kidney, liver, lung, mammary gland, placenta, small intestine, colon, spleen, prostate and testis. AKR1C3 catalyzes the conversion of aldehydes and ketones to alcohols. It catalyzes the reduction of prostaglandin (PG) D2, PGH2 and phenanthrenequinone (PQ) and the oxidation of 9-alpha,11-beta-PGF2 to PGD2, which functions as a bi-directional 3-alpha-, 17-beta- and 20-alpha HSD. It can interconvert active androgens, estrogens and progestins with their cognate inactive metabolites.

**Form:**PBS

**Molecular Weight:**37.8 kDa

**Sequences:**Met 1-Tyr323

**Purity:**> 95% by HPLC

**Concentration:**

**Endotoxin Level:**<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.