

**Recombinant Human USP5 / ISOT Protein (His tag)**

**Cat.NO.: TP05622**

3th Edition

**Synonyms:**ISOT

**Description:**Ubiquitin carboxyl-terminal hydrolase 5, also known as Deubiquitinating enzyme 5, Isopeptidase T, Ubiquitin thiolesterase 5, Ubiquitin-specific-processing protease 5, ISOT and USP5, is a member of the peptidase C19 family. USP5 contains 2 UBA domains and one UBP-type zinc finger. The UBP-type zinc finger domain interacts selectively with an unmodified C-terminus of the proximal ubiquitin. Both UBA domains are involved in polyubiquitin recognition. The UBP-type zinc finger domain crystallizes as a dimer linked by a disulfide bond between the Cys-195 residues of both molecules, but there is no evidence that the full-length USP5 exists as a dimer. USP5 cleaves linear and branched multiubiquitin polymers with a marked preference for branched polymers. USP5 is involved in unanchored 'Lys-48'-linked polyubiquitin disassembly. It binds linear and 'Lys-63'-linked polyubiquitin with a lower affinity. Knock-down of USP5 causes the accumulation of p53/TP53 and an increase in p53/TP53 transcriptional activity because the unanchored polyubiquitin that accumulates is able to compete with ubiquitinated p53/TP53 but not with MDM2 for proteasomal recognition.

**Form:**PBS

**Molecular Weight:**94.7 kDa

**Sequences:**Met 1-Ser 835

**Purity:**> 95% by HPLC

**Concentration:**

**Endotoxin Level:**<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.