
Recombinant Human APOM Protein (Fc tag)**Cat.NO.: TP05626**

3th Edition

Synonyms:apo-M;DADB-127H9.5;G3a;HSPC336;NG20

Description:ApoM (apolipoprotein M) is an apolipoprotein and member of the lipocalin protein family. The lipocalins share limited regions of sequence homology and a common tertiary structure architecture. They have an eight-stranded, antiparallel, symmetrical β -barrel fold, which is in essence a beta sheet which has been rolled into a cylindrical shape. Inside this barrel is located a ligand binding site. They transport small hydrophobic molecules such as steroids, bilins, retinoids, and lipids. Lipocalins have been associated with many biological processes, among them immune response, pheromone transport, biological prostaglandin synthesis, retinoid binding, and cancer cell interactions. Lipocalins are comparatively small in size, and are thus less complicated to study as opposed to large, bulky proteins. They can also bind to various ligands for different biological purposes. ApoM is associated with high density lipoproteins and to a lesser extent with low density lipoproteins and triglyceride-rich lipoproteins. ApoM is involved in lipid transport and can bind sphingosine-1-phosphate, myristic acid, palmitic acid and stearic acid, retinol, all-trans-retinoic acid and 9-cis-retinoic acid.

Form:PBS**Molecular Weight:**45.6 kDa**Sequences:**Met 1-Asn 188**Purity:**> 95% by HPLC**Concentration:****Endotoxin Level:**<1.0 EU per 1 ug of protein (determined by LAL method)**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.