

Recombinant Human Ganglioside GM2 activator Protein

Cat.NO.: TP05826

3th Edition

Synonyms:Ganglioside GM2 activator;Cerebroside sulfate activator protein;GM2-AP;Sphingolipid activator protein 3;SAP-3

Description:Ganglioside GM2 activator (GM2A) is a small glycolipid transport protein which acts as a substrate specific co-factor for the lysosomal enzyme β -hexosaminidase A (HEXB). HEXB together with GM2A, catalyzes the degradation of the ganglioside GM2, and other molecules containing terminal N-acetyl hexosamines. GM2A accommodate several single chain phospholipids and fatty acids, is a lipid transfer protein that stimulates the enzymatic processing of gangliosides, and also T-cell activation through lipid presentation. It extracts single GM2 molecules from membranes and presents them in soluble form to beta-hexosaminidase A for cleavage of N-acetyl-D-galactosamine and conversion to GM3.

Form:PBS

Molecular Weight:18.6 kDa

Sequences:Ser32-Ile193

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.