

**Recombinant Human Poliovirus Receptor-Related Protein 1/PVRL1 Protein(C-6His)**

**Cat.NO.: TP05898**

3th Edition

**Synonyms:** Poliovirus Receptor-Related Protein 1; Herpes Virus Entry Mediator C; Herpesvirus Entry Mediator C; HveC; Herpesvirus Ig-Like Receptor; HIgR; Nectin-1; CD111; PVRL1; HVEC; PRR1

**Description:** Nectin-1 is a type I transmembrane glycoprotein belonging to the Ig superfamily. Nectin-1 promotes cell-cell contacts by forming homophilic or heterophilic trans-dimers. Heterophilic interactions have been detected between Nectin-1 and Nectin-3 and between Nectin-1 and Nectin-4. Nectin ECDs contain three Ig like domains: an N terminal V type that mediates ligand binding, and two C2 type. Nectin-1 binds viral Glycoprotein D to mediate Herpesvirus (but not Poxvirus) entry into vaginal mucosa, sensory neurons and fibroblasts. In forming adherens junctions and synapses, Nectin-1 and Nectin-3 initiate cell-cell interactions, recruiting  $\alpha$ 5 $\beta$ 3 integrin extracellularly and cadherins intracellularly through afadin and other junctional proteins. These interactions organize the cytoskeleton, strengthen attachment to basement membrane and promote further cell-cell connections. Nectin-1 and Nectin-3 have been found to localize assymmetrically along the chemical synapse, with Nectin-1 primarily on the axonal side and Nectin-3 on the dendritic side. Deficiency of Nectin-1 can result in cleft lip/palate ectodermal dysplasia. Nectin-1 downregulation in epithelial cancers is mediated in part by ectodomain shedding, but it may contribute to invasiveness.

**Form:** PBS

**Molecular Weight:** 35.0 kDa

**Sequences:** Gln31-Thr334

**Purity:** > 95% by HPLC

**Concentration:**

**Endotoxin Level:** <1.0 EU per 1  $\mu$ g of protein (determined by LAL method)

**Storage:** Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.