

Recombinant Human Interleukin-5/IL-5 Protein(C-6His)

Cat.NO.: TP05940

3th Edition

Synonyms:Interleukin-5;IL-5;B-cell differentiation factor I;Eosinophil differentiation factor;T-cell replacing factor;TRF;IL5

Description:IL-5 is expressed in eosinophils, NK cells, TC2 CD8+ T cells, mast cells, CD45+ CD4+ T cells, gamma delta T cells and IL-1 beta activated endothelial cells. IL-5 acts as a growth and differentiation factor for both B cells and eosinophils. Relative to B cells, IL-5 appears to induce the differentiation of activated conventional B-2 cells into Ig-secreting cells. In addition, it induces the growth of B-1 progenitors as well as IgM production by B-1 cells.IL-5 appears to perform a number of functions on eosinophils. These include the down modulation of Mac-1,the upregulation of receptors for IgA and IgG,the stimulation of lipid mediator (leukotriene C4 and PAF) secretion and the induction of granule release.IL-5 also promotes the growth and differentiation of eosinophils.

Form:PBS

Molecular Weight:14.2 kDa

Sequences:Ile20-Ser134

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.