

**Recombinant Human Isocitrate Dehydrogenase 1/IDH1 Protein(C-8His)**

**Cat.NO.: TP05960**

---

3th Edition

**Synonyms:** Isocitrate Dehydrogenase [NADP] Cytoplasmic; IDH; Cytosolic NADP-Isocitrate Dehydrogenase; IDP; NADP(+)-Specific ICDH; Oxalosuccinate Decarboxylase; IDH1; PICD

**Description:** Isocitrate Dehydrogenase [NADP] Cytoplasmic (IDH1) belongs to the isocitrate and isopropylmalate dehydrogenases family. IDH1 exists as a homodimer, binding one magnesium or manganese ion per subunit. Mutations of IDH1 have been shown to cause metaphyseal chondromatosis with aciduria and are involved in the development of glioma. IDH1 plays a role in the regeneration of NADPH for intraperoxisomal reductions, such as the conversion of 2, 4-dienoyl-CoAs to 3-enoyl-CoAs, as well as in peroxisomal reactions that consume 2-oxoglutarate, namely the  $\alpha$ -hydroxylation of phytanic acid.

**Form:** PBS

**Molecular Weight:** 48.1 kDa

**Sequences:** Met 1-Leu414

**Purity:** > 95% by HPLC

**Concentration:**

**Endotoxin Level:** <1.0 EU per 1  $\mu$ g of protein (determined by LAL method)

**Storage:** Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.