

Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

Recombinant Human Mannoside Acetylglucosaminyltransferase 2/GlcNAc-TII Protein(C-6His)

Cat.NO.: TP06116

3th Edition

Synonyms:Alpha-1;6-Mannosyl-Glycoprotein 2-Beta-N-Acetylglucosaminyltransferase; Beta-1;2-N-acetylglucosaminyltransferase II; GlcNAc-T II; NT-II; Mannoside Acetylglucosaminyltransferase 2; N-Glycosyl-Oligosaccharide-Glycoprotein N-Acetylglucosaminyltransferase II; MGAT2

Description:Mannoside Acetylglucosaminyltransferase 2 (MGAT2) is a single-pass type II membrane protein that contains the typical glycosyltransferase domains: a short N-terminal cytoplasmic domain, a hydrophobic non-cleavable signal-anchor domain and a C-terminal catalytic domain. MGAT2 catalyzes an essential step in the conversion of oligo-mannose to complex N-glycans. Defects in MGAT2 are the cause of congenital disorder of glycosylation type 2A.

Form:PBS

Molecular Weight: 49.3 kDa

Sequences: Arg30-Gln447

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1