

Recombinant Mouse ICAM-2 / CD102 Protein (Fc tag)(Fc tag)

Cat.NO.: TP06238

3th Edition

Synonyms:CD102;Icam-2;Ly-60

Description:Intercellular adhesion molecule 2 (ICAM-2, CD102), belongs to the ICAM family consisting of three members identified as ligands for integrin receptors. It is a type I transmembrane glycoprotein with two Ig-like C2-type domains, and binds to the leukocyte integrins LFA-1 (CD11a/CD18) and Mac-1 (CD11b/CD18). As a second ligand of leukocyte function-associated antigen-1, ICAM-2 functions as a costimulatory molecule for effector cells. ICAM-2 is mainly expressed on vascular endothelial and hematopoietic cells. Interactions of ICAM-2 and the integrin receptors mediate cell adhesion in a wide range of lymphocyte, monocyte, natural killer cell, and granulocytewith other cells, and play important roles in many adhesion-dependent immune and inflammation responses, such as T cell aggregation, NK-cell cytotoxicity and migration, lymphocyte recirculation, etc. Serum levels of ICAM-2 correlated significantly with the inflammatory and course sequences of trichinosis in mice and had a similar relation with blood eosinophilia. So, estimation of ICAM-2 serum levels may prove useful in diagnosis of trichinosis recent infections, and in monitoring the prognosis and response to treatment.

Form:PBS

Molecular Weight:50 kDa

Sequences:Met 1-Gln 222

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.