

## Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

## Recombinant Rat VEGFR1 / FLT-1 Protein (His tag)

Cat.NO.: TP06244

3th Edition

Synonyms:FLT1;Flt-1;Vegfr1

**Description:** Vascular endothelial growth factor receptor 1, also known as VEGFR-1, Fms-like tyrosine kinase 1, Tyrosine-protein kinase FRT, Tyrosine-protein kinase receptor FLT, Vascular permeability factor receptor and FLT1, is a single-pass type I membrane protein and secreted protein which belongs to the protein kinase superfamily, Tyr protein kinase family and CSF-1/PDGF receptor subfamily. VEGFR-1 / FLT1 contains seven Iglike C2-type (immunoglobulin-like) domains and one protein kinase domain. VEGFR-1 / FLT1 is expressed mostly in normal lung, but also in placenta, liver, kidney, heart and brain tissues. It is specifically expressed in most of the vascular endothelial cells, and also expressed in peripheral blood monocytes. VEGFR-1 / FLT1 is not expressed in tumor cell lines. VEGFR-1 / FLT1 is an essential receptor tyrosine kinase that regulates mammalian vascular development and embryogenesis. EGF-induced angiogenesis requires inverse regulation of VEGFR-1 and VEGFR-2 in tumor-associated endothelial cells. VEGFR-1 / FLT1 is a receptor for VEGF, VEGFB and PGF. It has a tyrosine-protein kinase activity. The VEGF-kinase ligand/receptor signaling system plays a key role in vascular development and regulation of vascular permeability.Immune Checkpoint Immunotherapy Cancer Immunotherapy Targeted Therapy

Form:PBS

Molecular Weight:84 kDa

Sequences:Met1-Glu758

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1