

---

**Recombinant Human CD28 Protein (Fc tag)(Fc tag)**

**Cat.NO.: TP06513**

3th Edition

**Synonyms:** Tp44

**Description:** CD28 (Cluster of Differentiation 28) is a disulphide-bonded glycoprotein belonging to the immunoglobulin (Ig) superfamily, and structurally consists of a single Ig V-like extracellular domain, a transmembrane domain and an intracellular domain. Mouse CD28 is constitutively expressed on the surface of all murine T cells and on developing thymocytes as disulfide-linked homodimers or as monomers. CD28 can binds the B7-1 and B7-2 ligand, and together perform important functions in the T and B cell response pathways. B7/CD28 family members, which can augment or antagonize T-cell receptor signaling, in the regulation of central and peripheral T-cell tolerance. CD28 is thus involved in T-cell activation, the induction of cell proliferation and cytokine production and promotion of T-cell survival. Immune Checkpoint Immune Checkpoint Detection: Antibodies Immune Checkpoint Detection: ELISA Antibodies Immune Checkpoint Detection: IHC Antibodies Immune Checkpoint Detection: FCM Antibodies Immune Checkpoint Detection: WB Antibodies Immune Checkpoint Targets Co-stimulatory Immune Checkpoint Targets Immunotherapy Cancer Immunotherapy Targeted Therapy

**Form:** PBS

**Molecular Weight:** 42 kDa

**Sequences:** Met 1-Pro 152

**Purity:** > 95% by HPLC

**Concentration:**

**Endotoxin Level:** <1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:** Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.