

## Instruction manual FOR RESEARCH USE ONLY NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

## Recombinant Mouse SDF2 / SDF-2 Protein (His tag)

Cat.NO.: TP06783

3th Edition

Synonyms: Al853825

**Description:**Stromal derived factors (SDFs) are a loosely defined group of molecules that are generated by stromal cells. Two of the stromal derived factors, SDF-1 and SDF-4 belong to the chemokine family. Other SDFs, such as SDF-2 and SDF-5 are not well defined and their biological functions are less known. SDF-2 is first isolated from the mouse stromal cell line ST2 as a secretory protein. The amino acid sequence deduced from the murine clone and the human homolog are conserved more than 92 %, and the aa sequence of SDF-2 shows similarity to those of yeast dolichyl phosphate-D-mannose, protein mannosyltransferases. SDF-1 and its receptor are strongly indicated in the progression of various cancers including breast cancer. SDF-2, SDF2-L1, SDF-4, and SDF-5 are ubiquitously expressed in various cancer cell lines and SDF-2, SDF-4 and SDF-5 are expressed in mammary tissues. These SDFs have prognostic value and warrant further investigation in their biological functions and clinical value.

Form:PBS

Molecular Weight: 22.8 kDa

Sequences: Met 1-Leu 211

Purity:> 95% by HPLC

**Concentration:** 

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

**Storage:**Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.

1/1