

Recombinant Mouse FLRG / Fstl3 Protein (His tag)

Cat.NO.: TP07019

3th Edition

Synonyms:E030038F23Rik;Flrg

Description:Follistatin-like 3 (FLRG/Fstl3) is a secreted glycoprotein of the follistatin-module-protein family. It may have a role in leukemogenesis. FLRG/Fstl3 is a recently described member of the FST family having an overall structure and activity profile similar to that of FST, including binding and neutralization of activin. FLRG/Fstl3 is expressed in a wide range of adult tissues, not detected in hematopoietic cells except in patients with a B cell chronic leukemia and a translocation. Isoform 1 or the secreted form is a binding and antagonizing protein for members of the TGF-beta family, such as activin, BMP2 and MSTN. Inhibits activin A-, activin B-, BMP2- and MSDT-induced cellular signaling; more effective on activin A than on activin B. Involved in bone formation; inhibits osteoclast differentiation. Involved in hematopoiesis; involved in differentiation of hemopoietic progenitor cells, increases hematopoietic cell adhesion to fibronectin and seems to contribute to the adhesion of hematopoietic precursor cells to the bone marrow stroma. Isoform 2 of FLRG/Fstl3 or the nuclear form of FLRG/Fstl3 is probably involved in transcriptional regulation via interaction with MLLT10. Modulation of activin and other TGF β superfamily signaling is the primary mechanism of action for both follistatin (FS) and FS-like 3 (FSTL-3). FLRG/Fstl3 is likely to be a local regulator of activin action in gonadal development and gametogenesis and, further, that activin appears to have important actions in gonadal development and function that are critical for normal reproduction.

Form:PBS

Molecular Weight:26 kDa

Sequences:Met 1-Val 256

Purity:> 95% by HPLC

Concentration:

Endotoxin Level:<1.0 EU per 1 ug of protein (determined by LAL method)

Storage:Can be stored at +4°C short term (1-2 weeks). For long term storage, aliquot and store at -20°C or -70°C. Avoid repeated freezing and thawing cycles.